



1/10




FIG. 1



2/10




3/10



4/10


FIG. 5

#	cDNA
1	1A1
2	2B1 EST
3	2B1
4	2B2
5	3A1 h.
6	3A2
7	4A1
8	$\beta$ -actin
9	GAPDH
10	ext. std.



5/10

FIG. 6



6/10

FIG. 7a






FIG. 7b




7/10


*FIG. 8a*




*FIG. 8b*




*FIG. 8c*




*FIG. 8d*



*FIG. 8e*




*FIG. 8f*



*DISC-SHAPED PLATFORM*

*FIG. 8g*

*MICROTITER / NANOTITER-TYPE  
DEVICES (96/384/...)*



8/10

MM




FIG. 9c

REGENERATION  
50% UREA




FIG. 9b

PM










FIG. 9a




9/10

FIG. 10

		epi fluorescence			resonance		
	proteo-chips incubated with	image	spot means [counts]	signal/ noise*)	image	spot means [counts]	signal/ noise*)
I)	Cy5-labelled IL6 (500ng/ml, 10 microL, 2h incubation)		46	7.0		1100	69.2
II)	Cy5 labelled hCG (50ng/ml, 10 microL, 2h incubation)		32	5.0		646	75.1
III)	Cy5-labelled with mixture of anti-IL6 (polyclonal) /IL6 (50ng/ml, 10 microL, 12h incubation)		25	3.8		296	44.1

\*) noise [counts] defined as standard deviation of image background

\*) noise [counts] defined as standard deviation of image background



**Figure 12**

